




ENDING PREGNANCY WITH PILLS





For pregnancies up to 9 weeks



AM I PREGNANT?




I DON'T WANT TO BE PREGNANT!



CAN I TAKE PILLS TO END PREGNANCY?



WHAT ARE THE PILLS?



MISOPROSTOL:
DETAILS TO BE COMPLETED LOCALLY

WHAT SHOULD I DO BEFORE TAKING THE PILLS?


1




**BEFORE TAKING THIS
MEDICATION,
SEE A HEALTHCARE
PROVIDER IF YOU ...**

- have a bleeding disorder
- have severe anemia
- are allergic to the medications
- have a serious illness
- have an IUD - needs to be removed
- are more than 9 weeks pregnant
- have severe lower abdominal pain

2



3



HOW DO I TAKE THE PILLS?


3 DOSES

1ST DOSE



Put 4 pills under your tongue. Keep pills under your tongue until they melt, or for 30 minutes, then swallow whatever is left. (1 pill = 200 micrograms, 800 micrograms total)

Wait 3 hours then take 2nd dose.




2ND DOSE



Put 4 pills under your tongue. Keep pills under your tongue until they melt, or for 30 minutes, then swallow whatever is left. (1 pill = 200 micrograms, 800 micrograms total)

Wait 3 hours then take 3rd dose.




3RD DOSE




Put 4 pills under your tongue. Keep pills under your tongue until they melt, or for 30 minutes, then swallow whatever is left. (1 pill = 200 micrograms, 800 micrograms total)

WHAT WILL HAPPEN?



BLEEDING AND CRAMPING



FEELING BETTER

WHAT ARE THE POSSIBLE SIDE EFFECTS?


VOMITING / NAUSEA



HEADACHES



FEVER / CHILLS




DIARRHEA




GO TO THE HEALTH CLINIC IF YOU HAVE...




HEAVY BLEEDING FOR MORE THAN 2 HOURS IN A ROW
Immediately go to the health clinic



GO TO THE HEALTH CLINIC IF YOU HAVE...




SEVERE PAIN/SICKNESS FOR MORE THAN A DAY
Immediately go to the health clinic




GO TO THE HEALTH CLINIC IF SEVERE SIDE EFFECTS LAST MORE THAN A DAY



FEVER / CHILLS




DIARRHEA




GO TO THE HEALTH CLINIC IF SEVERE SIDE EFFECTS LAST MORE THAN A DAY




VOMITING / NAUSEA



HEADACHES





WILL I BE SUCCESSFUL IN ENDING MY PREGNANCY?



MOST WOMEN USING THESE PILLS TO END THE PREGNANCY HAVE A SUCCESSFUL ABORTION WITH NO PROBLEMS.

About 1 or 2 out of 10 women will need further care to end the pregnancy.

WHAT CAN I DO TO AVOID FUTURE PREGNANCY?





RETURN FOR YOUR FOLLOW-UP VISIT ON:

_____/_____/_____



QUESTIONS? CALL:



IN CASE OF EMERGENCY, GO TO THE NEAREST HOSPITAL
