


Print and Assembly Instructions for Gestational Dating Wheel for MA (both mife-miso and miso-only versions)

Note: It is very important that the small wheel be centered perfectly on the big wheel. If it is not, the number of days of gestation can be skewed and an inaccurate reading will occur. For this reason, it is a good idea to print and assemble a sample copy of the wheel before completing the entire print run.

1. The PDF is arranged with the front side of the big wheel as Page 1, the back side of the big wheel as Page 2, and the small wheel as Page 3.
2. To print the big wheel, ensure that you have selected the “two-sided printing” option on your computer. The PDFs are arranged so the wheels are centered on the page, and therefore your print-out should have the front and back sides of the wheel lined up evenly on opposite sides of the page. Print the desired quantity of Pages 1 and 2 together. Do not print page 3 at this time.
3. To print the small wheel, simply print the quantity of Page 3 that you need.
4. Neatly cut out both the large and small wheels along their outer lines – the PDFs have the correct size dimensions so the two wheels will fit together.
5. Center the small wheel over the side of the large wheel that shows the months and days of the year.
6. Use a brad, grommet, or similar connecting device to secure the two wheels together. The brad should anchor the wheels only at the center point, still allowing them to spin freely. Also, ensure that the connecting device does not obscure any text.
7. At this point, your Gestation Dating Wheel for MA should be ready to use. Please make sure that all text is legible and the small wheel is perfectly centered on the big wheel.

Black outline is trim line



First-trimester medical abortion — mifepristone and misoprostol

Instructions to use the wheel

- Place the arrow marked “First day of last menstruation” on the corresponding date.
- If today's date is inside the green zone you are at 9 weeks or less gestation and can begin taking mifepristone today and then misoprostol 1–2 days later.
- If today's date falls inside the yellow zone you are between 9 and 12 weeks gestation and you can use mifepristone and misoprostol only with medical supervision.
- If today's date is in the red zone you cannot use mifepristone and misoprostol.

Dosage and route of administration for green zone

- Swallow 200 mg of mifepristone.
- One or 2 days later, place 4 pills of 200 mcg each of misoprostol (800 mcg total) under the tongue or between the gum and the cheek of the mouth (2 pills on each side).
- After 30 minutes swallow anything remaining of the pills.

Warning Signs

Women should contact their health-care provider immediately if they experience:

- Excessive bleeding: soaking more than 2 sanitary pads per hour for 2 consecutive hours, especially if accompanied by prolonged dizziness, lightheadedness, and increasing fatigue.
- Fever that occurs any day after the day misoprostol is taken.
- Unusual or bad-smelling vaginal discharge, especially if accompanied by severe cramps or abdominal pain.
- Severe abdominal pain that occurs any day after the day misoprostol is taken.
- Feeling very sick with or without fever, and persistent, severe nausea or vomiting after the day misoprostol is used.

Black outline is trim line

